



04.1120.5017

MFM ANAPLAS BEEF MINERAL MEDICATED

For Further Manufacture of Feed

CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

For the control of bacterial pneumonia associated with shipping fever complex susceptible to chlortetracycline and control of active infection of anaplasmosis caused by *Anaplasma marginale* susceptible to chlortetracycline.

ACTIVE DRUG INGREDIENT

Chlortetracycline..... 2.0 g/lb

GUARANTEED ANALYSIS

Calcium, Min.....	14.50%
Calcium, Max.....	17.40%
Phosphorus, Min.....	8.00%
Salt, Min.....	13.70%
Salt, Max.....	16.30%
Magnesium, Min.....	1.00%
Potassium, Min.....	0.40%
Cobalt, Min (ppm).....	50
Copper, Min (ppm).....	2,500
Iodine, Min (ppm).....	200
Manganese, Min (ppm).....	5,000
Selenium, Min (ppm).....	26.4
Zinc, Min (ppm).....	4,000
Vitamin A, Min (IU/lb).....	300,000
Vitamin D3, Min(IU/lb).....	30,000
Vitamin E, Min(IU/lb).....	100

INGREDIENTS

Monocalcium phosphate, dicalcium phosphate, calcium carbonate, salt, processed grain by-products, molasses products, potassium sulfate, magnesium sulfate, yeast culture, manganese sulfate, zinc sulfate, copper sulfate, sodium selenite, ethylenediamine dihydroiodide, cobalt sulfate, vitamin A acetate, vitamin D3 supplement, vitamin E supplement, red iron oxide, artificial flavors.

FEEDING DIRECTIONS

For the control of bacterial pneumonia: Thoroughly mix 350 lbs. of **MFM Anaplas Beef Mineral** with 1,650 lbs. of non-medicated feed and feed the resulting medicated feed at a rate of 1 lbs. per head per day to deliver 350 mg CTC. Beef cattle should receive 350 mg CTC per head daily on a continuous basis for the stated drug claim.

Control of Anaplasmosis in Beef Cattle under 700 lbs: Thoroughly mix 350 lbs. of **MFM Anaplas Beef Mineral** with 1,650 lbs. of non-medicated feed and feed the resulting medicated feed at a rate of 1 lbs. per head per day to deliver 350 mg CTC. Beef cattle should receive 350 mg CTC per head daily on a continuous basis for the stated drug claim.

Control of Anaplasmosis in Beef Cattle under 700 lto 1,000 lbs.: Thoroughly mix 500 lbs. of **MFM Anaplas Beef Mineral** with 1,500 lbs. of non-medicated feed and feed the resulting medicated feed at a rate of 1 lb. per head daily to deliver 500 mg CTC. Beef cattle 700 to 1,000 lb. should receive 0.5 mg CTC per lb. of bodyweight daily on a continuous basis for the stated drug claim.

Warning: A withdrawal period has not been established for this product in pre-ruminating calves to be processed for veal.

Caution: Use as directed. Consumption of selenium should not exceed 3 mg per head per day.

Warning: This product which contains added copper, should not be fed to sheep or any species thata have a low tolerance to supplemental copper.

Manufactured for:
MARTINDALE FEED MILL
(A Division of Alan Ritchey, Inc)
Valley View, Texas

(800)366-9584
www.martindalefeed.com