



04.5501.5011 and 04.5501.2511

20-20 All Milk

CALF MILK REPLACER MEDICATED

For the control of coccidiosis caused by *Eimeria bovis* and *Eimeria zuernii* in replacement calves.

ACTIVE DRUG INGREDIENT

Lasalocid (as lasalocid sodium).....45 mg/pound
Equal to.....90 g/ton

* BOVATEC® brand of Lasalocid is a registered trademark of Alpha Inc.

GUARANTEED ANALYSIS

Crude Protein, Minimum.....20.00%
Crude Fat, Minimum.....20.00%
Crude Fiber, Maximum.....0.15%
Calcium, Minimum.....0.50%
Calcium, Maximum.....1.00%
Phosphorus, Minimum.....0.60%
Vitamin A, Minimum.....35,000 IU/lb
Vitamin D₃, Minimum.....7,500 IU/lb
Vitamin E, Minimum.....150 IU/lb

INGREDIENTS

Dried Whey, Dried Whey Product, Dried Milk Protein, Dried Skimmed Milk, Animal Fat preserved with BHA, Lecithin, Dicalcium Phosphate, Vitamin A Acetate, D-Activated Animal Sterol (source of Vitamin D₃), Vitamin E Supplement, Menadione Dimethylpyrimidinol Bisulfite (source of Vitamin K Activity), Riboflavin Supplement, Calcium Pantothenate, Niacin Supplement, Thiamine Mononitrate, Pyridoxine Hydrochloride, Choline Chloride, Vitamin B₁₂ Supplement, Biotin, Folic Acid, Ascorbic Acid, Magnesium Sulfate, Manganese Sulfate, Ferrous Sulfate, Zinc Sulfate, Cobalt Sulfate, Copper Sulfate, Calcium Iodate, Sodium Selenite, Sodium Silico Aluminate, Natural and Artificial Flavors.

Feeding Instructions on Back of Tag

WARNING: A withdrawal period has not been established for lasalocid in pre-ruminating calves.

DO NOT USE IN CALVES TO BE PROCESSED FOR VEAL.

CAUTION: The safety of lasalocid in unapproved species has not been established.

Manufactured For
MARTINDALE FEED MILL
Valleyview, TX 76272 U.S.A.

FEEDING PROGRAM

No whole milk is required after the colostrum feeding period when calf milk replacer is used.

MIXING AND FEEDING INSTRUCTIONS

Mix milk replacer powder at a rate of 2 cups (8 ounces) per 2 quarts of water (100° -120° F).

Feed approximately 2 quarts of reconstituted milk replacer TWICE A DAY.

Calves will vary in feed requirements depending on size and age.

*Measuring cup in each bag holds approx. 4 oz. dry weight.

FOR BEST RESULTS:

- Provide starter (grain mix) for calves when they are about 1 week of age and feed free choice up to 4 lbs. per day.
- For herd replacement calves, milk feeding can be discontinued when animals are about 6 to 7 weeks of age. However, a longer feeding period is recommended for small and poor performing calves.

GOOD MANAGEMENT PRACTICES:

- Good newborn care is crucial to maximize calf survival rates. Immediate feeding of colostrum is one of the most important steps to increase the survival and health of newborn calves. Feed 2 quarts of high quality, first milking colostrum within the first 30 minutes after birth. Feed an additional 2 quarts at 6 to 12 hours of age. Continue feeding colostrum through the third day after birth.
- Begin your milk replacer program on the 4th day. Feed twice daily, at regular intervals, according to label directions. Wash pails thoroughly after each usage.
- Have fresh, clean water available at all times. Dry hay can be offered after the first week.
- Calves should be housed in clean, dry individual pens that are well ventilated and draft free. Avoid sudden changes in temperature.

23253311
23253312

312B10B

Net. Weight on Bag

MMRC

(800)366-9584
www.martindalefeed.com