



04.1143.4017

## MFM BLOAT MINERAL MEDICATED

For Further Manufacture of Feed

For prevention of legume (alfalfa, clover) and wheat pasture bloat in cattle  
when fed continuously during exposure to bloat-producing conditions.

### ACTIVE DRUG INGREDIENT

Poloxalene ..... 4.4% (20g/lb)

### GUARANTEED ANALYSIS

Calcium, Min .....	8.20%
Calcium, Max .....	9.80%
Phosphorus, Min .....	3.00%
Salt, Min .....	22.00%
Salt, Max .....	26.40%
Magnesium, Min .....	4.00%
Cobalt, Min (ppm).....	50
Copper, Min (ppm) .....	625
Iodine, Min (ppm) .....	200
Manganese Min (ppm) .....	2,500
Selenium, Min (ppm) .....	8.80
Zinc, Min (ppm) .....	1,500
Vitamin A, Min (IU/lb) .....	150,000
Vitamin D3, Min (IU/lb) .....	15,000
Vitamin E, Min (IU/lb) .....	50

### INGREDIENTS

Salt, processed grain by-products, calcium carbonate, monocalcium phosphate, dicalcium phosphate, magnesium oxide, molasses products, yeast culture, manganese sulfate, zinc sulfate, copper sulfate, sodium selenite, ethylenediamine dihydroiodide, cobalt sulfate, vitamin A acetate, vitamin D3 supplement, vitamin E supplement, red iron oxide, and artificial flavoring.

### FEEDING DIRECTIONS

Thoroughly mix 500 lbs. of **MFM Bloat Mineral** mixed with 1,500 lbs. un-medicated feed, fed at a rate of 2 lbs. per head daily will deliver the recommended amount of poloxalene for a 1,000 lb. animal. Cattle should receive 1.0 to 2.0 g of poloxalene per 200 lbs. of bodyweight per day for the above drug claim. For optimum protection against bloat, begin feeding the medicated mixture at least 3 days prior to grazing bloat-producing pasture. Provide one covered mineral feeder for each 5—10 animals; and locate mineral feeders where animals congregate (watering and loafing areas).

It is essential that each animal consume the total recommended daily dosage of poloxalene for best bloat protection. If the average daily intake of the resulting medicated feed drops significantly below expected, be prepared immediately to relocate and increase average daily intake. If animals do not respond immediately with an increase in daily intake, remove the animals from bloat-producing pasture until the proper intake is re-established. Care should be taken to supply a clean, fresh source of water to all animals.

**CAUTION: Use as directed.** Consumption of selenium should not exceed 3 mg per head daily.

**Warning:** This product which contains added copper, should not be fed to sheep or any species that have a low tolerance to supplemental copper.

Manufactured For:

**MARTINDALE FEED MILL**

(A Division of Alan Ritchey, Inc)

Valley View, Texas 76272

**(800)366-9584**  
**www.martindalefeed.com**