



01.0020.5011

MFM 18% CALF STARTER (For Ruminants Only) -MEDICATED-

To be fed to calves approximately 3 days of age up to 3 months.

To increase rate of gain in pasture cattle (slaughter, stocker, feeder cattle, dairy or beef replacement heifers); for the control of coccidiosis caused by *Eimeria bovis* and *E. zuernii*.

ACTIVE DRUG INGREDIENT

Lasalocid (as lasalocid sodium)..... 35 mg/lb

CAUTION: The Safety of Lasalocid in unapproved species has not been established. Do not allow horses or other equine species access to lasalocid as ingestion may be fatal.

GUARANTEED ANALYSIS

Crude Protein, Min	18.00%
Crude Fat, Min.....	4.50%
Crude Fiber, Max.....	7.00%
Acid Deterent Fiber, Max.....	9.50%
Calcium, Min.....	0.80%
Calcium, Max.....	1.25%
Salt, Min	0.40%
Salt, Max	0.70%
Phosphorus, Min	0.55%
Potassium, Min.....	0.80%
Selenium, Min. (ppm)	0.30
Vitamin A, Min. (IU/LB).....	7,000

INGREDIENTS

Grain products, plant protein products, processed grain by-products, dried whey, cane molasses, yeast culture, calcium carbonate, monocalcium phosphate, dicalcium phosphate, salt, magnesium oxide, magnesium sulfate, potassium sulfate, sodium bicarbonate, vitamin A acetate, vitamin D3 supplement, vitamin E supplement, hydrolyzed animal fat (preserved with BHT), silicon dioxide, menadione sodium bisulfite complex (source of vitamin K activity), vitamin B-12 supplement, riboflavin supplement, d-calcium pantothenate, niacin supplement, choline chloride, thiamine mononitrate, pyridoxine hydrochloride, folic acid, ethoxyquin (a preservative), ascorbic acid, d-biotin, manganese sulfate, copper sulfate, cobalt sulfate, zinc sulfate, artificial flavor, sodium selenite, ethylenediamine dihydrochloride, ferrous carbonate, ferrous sulfate.

FEEDING DIRECTIONS

For increased rate of gain, feed continuously at a rate of no less than 60 mg (1 3/4 lbs) or more than 200 mg (5 3/4 lbs) of lasalocid/head/day. For control of coccidiosis in cattle up to 800 lbs. of body weight, feed continuously to provide 1 mg per 2.2 lbs of body weight/head/day, not to exceed 360 mg (10 1/4 lbs)/head/day. Provide a source of fresh, clean water at all times. To insure freshness, offer only the amount calves will consume in one day. Clean feeders daily. Store feed in a dry, well ventilated area. Consult your veterinarian for a planned herd health program.

◆ DO NOT FEED MOLDY OR INSECT INFESTED FEED. ◆

WARNING: A withdrawal period has not been established for this product in pre-ruminating cattle.

Do not use in calves to be processed for veal.

CAUTION: Feed as directed. For ruminant animals only.

Manufactured By:

MARTINDALE FEED MILL

(a division of Alan Ritchey, Inc.)

Valley View, Texas 76272

Net Weight 50 lbs. (22.68 kg)

(800)366-9584
www.martindalefeed.com