



04-3313-9620

# MFM PASTURE-MAX 24% TUB



04-3313-9620

## PASTURE MAX 24% Supplement Block

For Mature Beef Cattle on Pasture

**GUARANTEED ANALYSIS:**

Crude Protein (min).....	24.00%
(Includes not more than 16.00% equivalent crude protein from non-protein nitrogen)	
Crude Fat (min).....	5.00%
Crude Fiber (max).....	6.00%
Calcium (min).....	0.25%
Calcium (max).....	0.75%
Phosphorus (min).....	0.25%
Salt (min).....	2.50%
Salt (max).....	3.00%
Magnesium (min).....	3.00%
Copper (min).....	45 ppm
Manganese (min).....	170 ppm
Selenium (min).....	3.3 ppm
Zinc (min).....	130 ppm
Vitamin A (min).....	15,000 IU/lb
Vitamin D-3 (min).....	1,500 IU/lb
Vitamin E (min).....	1.5 IU/lb

**INGREDIENT STATEMENT:**

Molasses Products, Processed Grain By-Products, Magnesium Oxide, Urea, Salt, Vegetable Oil, Monocalcium Phosphate, Dicalcium Phosphate, Calcium Propionate (a preservative), Roughage Products, Sodium Selenite, Calcium Carbonate, Manganese Sulfate, Zinc Sulfate, Copper Sulfate, Manganese Oxide, Zinc Oxide, Mineral Oil, Vitamin A Supplement, Vitamin D-3 Supplement, Vitamin E Supplement, Ethylenediamine Dihydroiodide, Calcium Iodate and Cobalt Carbonate.

**INTRODUCTORY PERIOD:** Following a 7 to 10 day acclimation period, average consumption should be 1 to 2 pounds per head per day for mature cattle with unrestricted forage supplies. Position the blocks 50 to 75 feet from loafing, feeding and watering areas. Some repositioning of blocks may be necessary as animals become acclimated to the presence of the blocks. Move blocks closer to increase consumption; further away to decrease consumption. If over-consumption occurs for more than two weeks and repositioning of blocks does not correct the situation, remove blocks and evaluate the overall feeding program. **DO NOT ALLOW STARVED ANIMALS FREE ACCESS TO THIS BLOCK.**

**NET WEIGHT: 200 LBS. (90.7 KG)**

**CAUTION:** Use as directed. Consumption of selenium should not exceed 3 mg per head daily.

**WARNING:** This product, which contains added copper, should not be fed to sheep or any species that have a low tolerance to supplemental copper.

**FEEDING DIRECTIONS**

1. Provide 1 block per 15 to 30 head. All animals should have equal access to blocks. Timid and smaller animals need their share.
2. Place blocks where animals congregate. Locate blocks throughout the pasture at loafing, grazing, feeding and watering areas. Adequate distance must be maintained between blocks to minimize crowding - 10 feet minimum.
3. Feed blocks continuously along with a plentiful source of forage and clean, fresh water. Animals should not run out of blocks. When a block is two-thirds consumed, provide a new block near it.
4. Cattle consume 1 to 2 lbs. per head daily based on a 1,000-lb. animal.
5. Consumption of this block may vary depending upon animal body condition, quality and quantity of forages, seasonal weather conditions, and most importantly, feeding locations of blocks with respect to loafing, grazing, feeding and watering areas.
6. If additional mineral and vitamin fortification is desired, provide an appropriate free choice mineral supplement in addition to this supplement.

For additional information, contact your Martindale Feed Mill representative.

Manufactured for:  
**MARTINDALE FEED MILL**  
(A division of Alan Ritchey, Inc.)  
Valley View, TX 76272

# 11146

1121