

04-7102-9620

MFM PASTURE MAX 24% COOKED TUB with ALTOSID® IGR.0025

2

Λ 5.

6

7.

8

9

Pasture Max 24% Cooked Tub with Altosid[®] IGR.0025

Prevents adult horn fly emergence from manure of treated cattle. ent Containing (S)-Meth ene* Insect Growth Regulator for Conti ous Feeding to

Cattle through the Horn Fly Season.	
ACTIVE INGREDIENT:	

ACTIVE INGREDIENT:	
(S)-Methoprene (CAS #65733-16-6)	
OTHER INGREDIENTS	
TOTAL	

* As the Insect Growth Regulator ALTOSID* GUARANTEED ANALYSIS

Crude Protein , (Min)	24.00%
Crude Fat (Min)	5.00%
Crude Fiber (Max)	2.50%
Calcium (Min)	1.00%
Calcium (Max)	1.50%
Phosphorus (Min)	1.00%
Salt (Min)	none added
Magnesium (Min)	
Potassium (Min)	2.00%
Cobalt (Min)	9 ppm
Copper (Min)	
lodine (Min)	
Mangapese (Min)	450 ppm
Selenium (Min)	6 ppm
Zinc (Min)	mag 008
Vitamin A (Min)	
Vitamin D. (Min)	
Vitamin E (Min)	

FEED INGREDIENTS

Molasses products, animal protein products, processed grain by-products, monocalcium phosphate, dicalcium phosphate, magnesium oxide, vegetable oil, calcium carbonate, plant protein products, roughage products, copper sulfate, zinc oxide, manganese sulfate, sodium selenite, ethylenediamine dhydroiodide, zinc sulfate, manganous oxide, vitamin E supplement, vitamin A supplement, vitamin D-3 supplement, mineral oil, cobalt carbonate, calcium iodate.

Keep Out of Reach of Children

This product contains ethoxyquin a preservative. Altosid[®] is a registered trad mark of Wellmark International EPA Reg No. 2724-787-7698 EPA Est. No. 7698-TX-001

NET WEIGHT: 200 LB. (90.7 KG)

Manufactured by: Ridley USA Inc., 111 W. Cherry Street, Suite 500, Mankato, Mn 56001

ENVIRONMENTAL HAZARDS: Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

GENERAL INFORMATION

This pesticide product is to be used as a cattle feed-through pesticide for the control of horn flies. All feed ingredients listed or this product label must adhere to the specifications and limits (if applicable) ascribed by the Food and Drug Administration (FDA) for cattle feed use when mixed with this pesticide product. This product contains ethoxyquin, a preservative. The amount of ethoxyquin may vary according to the individual manufacturer's formulation and will not exceed the 150 ppm maximum limit established for ethoxyquin by FDA, per 21 CFR 573.380. FEEDING AND MANAGEMENT INSTRUCTIONS:

Feed this product to cattle only. When used as directed, MFM Pasture Max 24% Cooked Tub with Attosid⁶ IGR.0025 will provide sufficient (S)-methoprene insect growth regulator to prevent the emergence of adult horn flies from manure of treated cattle. Existing adult horn flies will not be affected. Start feeding before horn flies appear and continue use until cold weather marks the end of the horn fly season. Allow free choice to cattle. Cattle should consume 34 mg of (s)-methoprene per 100 lbs of cattle body weight per month to achieve horn fly control. This is equivalent to an average of 1.6 ounces of this product per 100 lbs of cattle body weight per day, which is equivalent to 1 pound (16 ounces) per day for a 1000-lb. cow. To open: Remove protective cover

- Assure cattle are not starved for nutrients prior to feeding MFM Pasture Max 24% Cooked Tub with Altosid® IGR.0025.
- This product contains added copper. Do not feed to cattle in close proximity to sheep or other species that have a low tolerance to supplemental copper. Remove all other self-limiting nutritional supplements while cattle are consuming this product.
- Offer cattle one MFM Pasture Max 24% Cooked Tub with Altosid® IGR.0025 for every 15 to 25 head of cattle. Locate MFM Pasture Max 24% Cooked Tub with Altosid® IGR.0025 where cattle congregate (watering, loafing, shade areas).
- Replenish MFM Pasture Max 24% Cooked Tub with Altosid® IGR.0025 on a regular basis as needed.
- If intake of MFM Pasture Max 24% Cooked Tub with Altosid® IGR.0025 is below 1.6 oz. per 100 lbs. cattle body weight per day, increase the number of feeding locations and/or change locations to areas more frequented by cattle
- If intake of MFM Pasture Max 24% Cooked Tub with Altosid® IGR.0025 is above 1.6 oz. per 100 lbs. cattle body weight per day, reduce the number of feeding locations and/or change locations to areas less frequented by cattle
- Introduction of this product after adult horn fly infestation is established will require treatment of cattle with adulticides if elimination of the adult fly population is desired.

STORAGE AND DISPOSAL: Do not contaminate water, food or feed by storage or disposal. STORAGE: STORAGE AND DISPOSAL: Up not contaminate water, lood or feed by storage or disposal. STORAGE Store in a cool, dry place. Do not stack anything on top of this container without placing a silp sheet or wooden pallet between containers. PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. Triple rinse (or equivalent). Then offer for recycling if available or puncture and dispose of in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke. Batch # / MEG

72031

