



01-0075-5014

Safe-Guard® Cattle Dewormer Cubes 0.100%

Medicated

A Dewormer for Beef Cattle on Pasture and Non-Lactating Dairy Cattle.

For the treatment and control of: **Lungworms:** Adult *Dictyocaulus viviparus*; **Stomach worms:** Adult brown stomach worms (*Ostertagia ostertagi*), Adult and fourth stage larvae barberpole worms (*Haemonchus contortus*), fourth stage larvae barberpole worms (*H. placei*), and Adult and fourth stage larvae small stomach worms (*Trichostrongylus axei*); **Intestinal worms:** (Adult and fourth stage larvae): hookworms (*Bunostomum phlebotomum*), thread-necked intestinal worms (*Nematodirus helvetianus*), small intestinal worms (*Cooperia punctata* & *C. oncophora*), bankrupt worms (*Trichostrongylus colubriformis*), and nodular worms (*Oesophagostomum radiatum*).

-ACTIVE DRUG INGREDIENT-

Fenbendazole..... 0.100%

-GUARANTEED ANALYSIS-

Crude Protein, Min.....	20.00%
Crude Fat, Min.....	2.50%
Crude Fiber, Max.....	13.00%
Acid Detergent Fiber, Max.....	15.00%
Calcium, Min.....	1.50%
Calcium, Max.....	2.50%
Phosphorus, Min.....	0.60%
Salt, Min.....	0.50%
Salt, Max.....	1.00%
Potassium, Min.....	1.00%
Selenium, Min (ppm).....	0.20
Vitamin A Min (IU/LB).....	10,000

-INGREDIENTS-

Processed grain by-products, plant protein products, grain products, forage products, cane molasses, calcium carbonate, salt, vitamin A acetate, D-activated animal sterol (source of vitamin D3), vitamin E supplement, sodium selenite, manganese sulfate, zinc sulfate, ferrous sulfate, copper sulfate, ethylenediamine dihydriodide, cobalt carbonate.

-FEEDING DIRECTIONS-

Feed as the sole ration for One (1) day. Feed at a rate of .5 pounds per 100 pounds of body weight. (Example: 1,000 lb. cow, feed 5 lbs.) This will provide 2.27 mg of fenbendazole per pound of body weight.

Do not underdose. Ensure each animal receives a complete dose based on a current body weight. Underdosing may result in ineffective treatment and encourage the development of parasite resistance.

WITHDRAWAL PERIODS AND RESIDUE WARNINGS

Milk taken during treatment and for 60 hours after the last treatment must not be used for human consumption. Cattle must not be slaughtered for human consumption within 13 days following last treatment with this drug product. Not for use in beef calves less than 2 months of age, dairy calves, and veal calves. A withdrawal period has not been established for this product in pre-ruminating calves.

OTHER WARNINGS

Parasite resistance may develop to any dewormer, and has been reported for most classes of dewormers. Treatment with a dewormer used in conjunction with parasite management practices appropriate to the geographic area and the animal(s) to be treated may slow the development of parasite resistance. Fecal examinations or other diagnostic tests and parasite management history should be used to determine if the product is appropriate for the herd, prior to the use of any dewormer. Following the use of any dewormer, effectiveness of treatment should be monitored (for example, with the use of a fecal egg count reduction test or another appropriate method). A decrease in a drug's effectiveness over time as calculated by fecal egg count reduction tests may indicate the development of resistance to the dewormer administered. Your parasite management plan should be adjusted accordingly based on regular monitoring.

Consult your veterinarian for assistance in diagnosis, treatment and control of parasitism.
Approved by FDA under NADA # 131-675

*Safe-Guard® is a registered trademark of Intervet International B.V.

Manufactured By:
MARTINDALE FEED MILL
(a division of Alan Ritchey, Inc.)
Valley View, Texas 76272

(800)366-9584
www.martindalefeed.com