



04-1142-5017

# MFM PASTURE MAX GRAZER R (RUMENSIN) MINERAL -MEDICATED-

For Further Manufacture of Feed

For increased rate of weight gain in growing cattle on pasture or in a dry lot (stocker and feeder; dairy and beef replacement heifers).

### -ACTIVE DRUG INGREDIENT-

Monensin..... 1,600 g/ton

### -GUARANTEED ANALYSIS-

Calcium, Min.....	12.60%
Calcium, Max.....	15.10%
Phosphorus, Min .....	3.50%
Salt, Min .....	14.60%
Salt, Max .....	17.50%
Magnesium, Min .....	0.60%
Cobalt, Min (ppm).....	50
Copper, Min (ppm) .....	830
Iodine, Min (ppm) .....	200
Manganese, Min (ppm) .....	3,000
Selenium, Min (ppm) .....	26.40
Zinc, Min (ppm) .....	2,000.00
Vitamin A, Min (IU/lb) .....	300,000
Vitamin D3, Min (IU/lb) .....	30,000
Vitamin E, Min (IU/lb) .....	100

### -INGREDIENTS-

Processed grain by-products, calcium carbonate, salt, monocalcium phosphate, dicalcium phosphate, molasses products, potassium sulfate, magnesium sulfate, yeast culture, manganese sulfate, zinc sulfate, copper sulfate, ethylenediamine dihydroiodide, cobalt sulfate, vitamin A acetate, vitamin D<sub>3</sub> supplement, vitamin E supplement, mineral oil, red iron oxide and artificial flavoring.

### -FEEDING DIRECTIONS-

Thou roughly mix 500 lbs. of **MFM Grazer R (Rumensin)** Mineral with 1,500 lbs. of non-medicated feed. Feed one pound of this medicated mixture to provide 200 mg of monensin. Cattle should receive not less than 50 nor more than 200 mg monensin per head per day in not less than 1 lb. of feed or after the 5th day at the rate of 400 mg per head per day every other day in not less than 2 lbs. of feed for the stated drug claim. During the first 5 days of feeding the cattle should receive no more than 100 mg per day contained in at least 1 lb. of feed. Thoroughly mix 250 lbs. of MFM Grazer R (Rumensin) Mineral with 1,750 lbs. of a non-medicated feed. Feed one pound of this medicated mixture to provide 100 mg monensin, For additional informayion contact your MFM representative.

**CAUTION:** Do not allow horses or other equines access to feed containing monensin. Ingestion of monensin by equines has been fatal. Monensin-medicated cattle and goat feeds are safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions. Do not use undiluted. Feeding undiluted or mixing errors resulting in high concentrations of monensin has been fatal to cattle and could be fatal to goats. Do not exceed the levels of monensin recommended in the feeding directions, as reduced average daily gains may result. If feed refusals containing monensin are fed to other groups of cattle, the concentration of monensin in the refusals fed should be taken into consideration to prevent overdosing. Consumption of selenium should not exceed 3 mg per head daily.

**WARNING:** A withdrawal period has not been established for pre-ruminating calves. Do not use in calves to be processed for veal. This product, which contains added copper, should not be fed to sheep or species that have a low tolerance to supplemental copper.

Manufactured For:  
**MARTINDALE FEED MILL**  
(A Division of Alan Ritchey, Inc)  
Valley View, Texas 76272

**(800)366-9584**  
**www.martindalefeed.com**