



04.3310..3320

SWEETLIX PEST-A-SIDE/RABON BLOCK



Pest-A-Side Pressed Block with Rabon®

A Free Choice Supplement Block for Cattle



For control of fecal flies in manure of treated cattle. Prevents development of face flies, horn flies, house flies and stable flies in the manure of treated cattle.

Active Ingredient:

Tetrachloro-1-(2,4,5-Trichlorophenyl) vinyl dimethyl phosphate*	0.30%
Other Ingredients**	99.70%
	100.00%

* Each lb. contains 1.36 g Rabon®
** Refers only to ingredients which are not larvicidal

Guaranteed Analysis

Crude Protein (Min)	8.00%
Crude Fat (Min)	0.50%
Crude Fiber (Max)	7.00%
Calcium (Min)	7.00%
Calcium (Max)	8.00%
Phosphorus (Min)	2.00%
Salt (Min)	10.00%
Salt (Max)	12.00%
Magnesium (Min)	3.90%
Potassium (Min)	1.00%
Cobalt (Min)	60 ppm
Copper (Min)	180 ppm
Iodine (Min)	60 ppm
Manganese (Min)	500 ppm
Selenium (Min)	12 ppm
Zinc (Min)	700 ppm
Vitamin A (Min)	40,000 IU/lb.
Vitamin D-3 (Min)	5,000 IU/lb.
Vitamin E (Min)	20 IU/lb.

Feed Ingredients

Cane Molasses, Dehydrated Alkali, Cottonseed Meal, Lignite Sulfonate, Dolomitic Limestone, Magnesium Nitrate, Monocalcium Phosphate, Dicalcium Phosphate, Sulfur, Calcium Benzoate, Vitamin A Supplement, Vitamin D-3 Supplement, Vitamin E Supplement, Ferrous Sulfate, Manganese Oxide, Manganese Sulfate, Zinc Oxide, Zinc Sulfate, Calcium Iodate, Copper Sulfate, Cobalt Sulfate, Sodium Selenite, Plant Protein Products and Calcium Carbonate.

Keep Out of Reach of Children

CAUTION

EPA Reg. No. 73600-5 * EPA Est. No. 73600-TX-01

Manufactured By Sweetlix, PO Box 8500, Markato, MN 56002

Contact us at: 1-877-SWEETLIX * www.sweetlix.com

First Aid	
IF SWALLOWED	Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor.
IF ON SKIN	Take off contaminated clothing. Rinse skin with plenty of water for 15-20 minutes. Call poison control center or doctor for treatment advice.
IF IN EYES	Hold eyes open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or physician for treatment advice.

HOT LINE NUMBER: In case of emergency, contact Sweetlix (800) 325-1486.

NOTE TO PHYSICIANS AND VETERINARIANS: Symptomatology: Signs of organophosphate toxicity are those of cholinergic over stimulation: hypersalivation, excessive urination, diarrhea and dyspnea (respiratory distress, collapse and death from respiratory muscle paralysis). Treatment: Withdraw product immediately. Clinical includes 3 categories: muscarinic blocking agents (atropine sulfate), (2) cholinesterase reactivators (atropine and 2-PAM, pralidoxime chloride) and (3) emetics (i.e. syrup of ipecac), cathartics (i.e. vegetable oil) and absorbents (i.e. charcoal slurry).

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS:

CAUTION

Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Prolonged or frequently repeated skin contact may cause allergic reaction to some individuals.

USER SAFETY RECOMMENDATIONS

PERSONAL PROTECTIVE EQUIPMENT (PPE): Long-sleeved shirt and long pants, shoes and socks, chemical resistant gloves. Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing. Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Follow manufacturers instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENVIRONMENTAL HAZARDS: This compound is toxic to fish.

Rabon® is a registered trademark of KMG Chemicals, Houston, TX

NET WEIGHT: 33.3 LB. (15.10 KG)



DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

GENERAL INSTRUCTIONS

1. Feed 1 block per 10-15 head of cattle. This allows all animals equal access to blocks. Feed blocks at the rate of 0.88 oz. per 100 lb. of bodyweight per day. This intake will supply 0.07g. of larvicide per 100 lb. of bodyweight per day (2.1 grams per 100 lbs. bodyweight per month).
2. Place blocks where animals congregate. Locate blocks throughout pasture, placing them near water and feeding areas. Maintain adequate distance between blocks (10 ft. minimum).
3. Do not allow animals to run out of blocks. When a block is half-consumed, place a new block near it. Place very small pieces in troughs or feed pans.
4. Animals should consume about 8.8 oz. per head daily. When animals consume the block at the recommended rate, it will supply 3 mg of selenium.
5. Blocks can be fed to animals in confinement. Exact location of blocks to obtain desired consumption will vary between confinement feeding programs.
6. Feed as the only FREE CHOICE source of salt, other minerals and vitamins.
7. SWEETLIX® Pest-A-Side Pressed Block with Rabon® prevents the development of fecal flies in the manure of treated cattle, but it is not effective against existing adult flies. It may be necessary to use other control measures to reduce the population of existing adult flies. Supplemental fly control measures may be needed in and around buildings to control adult house flies and stable flies which can breed not only in animal manure but also in other decaying matter on the premises. In order to achieve optimum fly control, SWEETLIX® Pest-A-Side Pressed Block with Rabon® should be used in conjunction with other good management and sanitation practices. Start feeding SWEETLIX® Pest-A-Side Pressed Block with Rabon® early in the spring before flies begin to appear and continue feeding throughout the summer and into the fall. SWEETLIX® Pest-A-Side Pressed Block with Rabon® may be fed to cattle up to slaughter.
8. Follow feeding directions. Consumption of selenium should not exceed 3 mg per head daily.

STORAGE AND DISPOSAL: Do not contaminate water, food, or feed by storage or disposal.

Disposal: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or if allowed by State and Local Authorities, by burning. If burned, stay out of smoke. Wastes resulting from use of this product may be disposed of on site or at an approved waste disposal facility. For additional information, please contact your SWEETLIX representative.

03/05

(800)366-9584
www.martindalefeed.com